

Comment on
“Solutions of the Duffin-Kemmer-Petiau equation for a
pseudoscalar potential step in (1+1) dimensions”

T.R. Cardoso, L.B. Castro¹, and A.S. de Castro²

UNESP - Campus de Guaratinguetá
Departamento de Física e Química
12516-410 Guaratinguetá SP - Brazil

¹benito@feg.unesp.br

²castro@pq.cnpq.br.

Abstract

It is shown that the paper “Solutions of the Duffin-Kemmer-Petiau equation for a pseudoscalar potential step in (1+1) dimensions” by Abdelmalek Boumali has a number of misconceptions.

In a recent paper published in this Journal, Boumali [1] reports on solutions of the Duffin-Kemmer-Petiau (DKP) equation and draws different conclusions about Klein's paradox for spin-0 and spin-1 bosons. The purpose of this comment is to point that Ref. [1] has a number of misconceptions endangering its main conclusions.

The DKP equation for a free boson is given by [2]-[4]

$$(i\beta^\mu\partial_\mu - m)\Psi = 0 \quad (1)$$

where the matrices β^μ satisfy the algebra

$$\beta^\mu\beta^\nu\beta^\lambda + \beta^\lambda\beta^\nu\beta^\mu = g^{\mu\nu}\beta^\lambda + g^{\lambda\nu}\beta^\mu \quad (2)$$

and the metric tensor is $g^{\mu\nu} = \text{diag}(1, -1, -1, -1)$. A well-known conserved four-current is given by

$$J^\mu = \bar{\Psi}\beta^\mu\Psi \quad (3)$$

where the adjoint spinor $\bar{\Psi} = \Psi^\dagger\eta^0$, with $\eta^0 = 2\beta^0\beta^0 - 1$ in such a way that $(\eta^0\beta^\mu)^\dagger = \eta^0\beta^\mu$ (the matrices β^μ are Hermitian with respect to η^0). With the introduction of interactions, the DKP equation can be written as

$$(i\beta^\mu\partial_\mu - m - U)\Psi = 0 \quad (4)$$

where the more general potential matrix U is written in terms of 25 (100) linearly independent matrices pertinent to the five(ten)-dimensional irreducible representation associated to the scalar (vector) sector. In the presence of interactions J^μ satisfies the equation

$$\partial_\mu J^\mu + i\bar{\Psi}(U - \eta^0 U^\dagger \eta^0)\Psi = 0 \quad (5)$$

Thus, if U is Hermitian with respect to η^0 then four-current will be conserved. The potential matrix U can be written in terms of well-defined Lorentz structures. For the spin-0 sector there are two scalar, two vector and two tensor terms [5], whereas for the spin-1 sector there are two scalar, two vector, a pseudoscalar, two pseudovector and eight tensor terms [6]. Restricting ourselves to the spin-0 sector of the DKP theory and considering only scalar and vector terms, U is in the form

$$U = S^{(1)} + PS^{(2)} + \beta^\mu V_\mu^{(1)} + i[P, \beta^\mu]V_\mu^{(2)} \quad (6)$$

where P is a projection operator ($P^2 = P$ and $P^\dagger = P$) that picks out the component of the DKP spinor which satisfies the free Klein-Gordon equation. Note that this matrix potential leads to a conserved four-current but the same does not happen if, instead of

$i[P, \beta^\mu]$ one uses $\beta^\mu P$. With the representation for the β^μ matrices given by [7] (apparently the same representation as that one used in Ref. [1])

$$\beta^0 = \begin{pmatrix} \theta & \bar{\mathbf{0}} \\ \bar{\theta}^T & \mathbf{0} \end{pmatrix}, \quad \beta^i = \begin{pmatrix} \tilde{\mathbf{0}} & \rho_i \\ -\rho_i^T & \mathbf{0} \end{pmatrix}, \quad i = 1, 2, 3 \quad (7)$$

where

$$\begin{aligned} \theta &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & \rho_1 &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ \rho_2 &= \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \rho_3 &= \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix} \end{aligned} \quad (8)$$

$\bar{\mathbf{0}}$, $\tilde{\mathbf{0}}$ and $\mathbf{0}$ are 2×3 , 2×2 and 3×3 zero matrices, respectively, while the superscript T designates matrix transposition, the projection operator can be written as [5]

$$P = \frac{1}{3} (\beta^\mu \beta_\mu - 1) = -\beta^0 \beta^0 \beta^1 \beta^1 \beta^2 \beta^2 \beta^3 \beta^3 = \text{diag}(1, 0, 0, 0, 0) \quad (9)$$

The five-component spinor can be written as $\Psi^T = (\Psi_1, \dots, \Psi_5)$ in such a way that the DKP equation for a boson constrained to move along the X -axis decomposes into

$$\begin{aligned} D_0^{(+)} \Psi_1 &= -i(m + S^{(1)}) \Psi_2, & D_1^{(+)} \Psi_1 &= -i(m + S^{(1)}) \Psi_3 \\ D_0^{(-)} \Psi_2 - D_1^{(-)} \Psi_3 &= -i(m + S^{(1)} + S^{(2)}) \Psi_1 \\ \Psi_4 &= \Psi_5 = 0 \end{aligned} \quad (10)$$

where

$$D_\mu^{(\pm)} = \partial_\mu + iV_\mu^{(1)} \pm V_\mu^{(2)} \quad (11)$$

In this case J^μ decomposes into

$$J^0 = 2\text{Re}(\Psi_1 \Psi_2^*), \quad J^1 = -2\text{Re}(\Psi_1 \Psi_3^*), \quad J^2 = J^3 = 0 \quad (12)$$

If the terms in the potential matrix U are time-independent, one can write $\Psi(x, t) = \psi(x) \exp(-iEt)$ in such a way that the time-independent DKP equation decomposes into

$$(m + S^{(1)}) \frac{d}{dx} \left(\frac{1}{m + S^{(1)}} \frac{d\psi_1}{dx} \right) + 2iV_1^{(1)} \frac{d\psi_1}{dx} + k^2 \psi_1 = 0$$

$$\begin{aligned}\psi_2 &= \frac{1}{m + S^{(1)}} \left(E - V_0^{(1)} + iV_0^{(2)} \right) \psi_1 \\ \psi_3 &= \frac{i}{m + S^{(1)}} \left(\frac{d}{dx} + iV_1^{(1)} + V_1^{(2)} \right) \psi_1\end{aligned}\tag{13}$$

where

$$\begin{aligned}k^2 &= \left(E - V_0^{(1)} \right)^2 - \left(V_1^{(1)} \right)^2 + i \frac{dV_1^{(1)}}{dx} + \left(V_0^{(2)} \right)^2 - \left(V_1^{(2)} \right)^2 + \frac{dV_1^{(2)}}{dx} \\ &\quad - \left(m + S^{(1)} \right) \left(m + S^{(1)} + S^{(2)} \right) - \frac{iV_1^{(1)} + V_1^{(2)}}{m + S^{(1)}} \frac{dS^{(1)}}{dx}\end{aligned}\tag{14}$$

For this time-independent problem, J^μ has the components

$$J^0 = 2 \frac{E - V_0^{(1)}}{m + S^{(1)}} |\psi_1|^2, \quad J^1 = 2 \frac{V_1^{(1)} |\psi_1|^2 + \text{Im} \left(\frac{d\psi_1}{dx} \psi_1^* \right)}{m + S^{(1)}}\tag{15}$$

J^μ is not time dependent, so that ψ describes a stationary state.

The form $\partial_1 + iV_1^{(1)}$ in Eq. (10) suggests that the space component of the minimal vector potential can be gauged away by defining a new spinor

$$\tilde{\Psi}(x, t) = \exp \left[i \int^x d\zeta V_1^{(1)}(\zeta, t) \right] \Psi(x, t)\tag{16}$$

even if $V_1^{(1)}$ is time dependent. Furthermore, the elimination of the first derivative of a second-order differential equation, such as the term containing $V_1^{(1)}$ in (13), is a well-known trick in mathematics. Nevertheless, it seems that there is no chance to get rid from this term, except of course on condition that one imposes $V_1^{(2)} = 0$.

The DKP equation is invariant under the parity operation, i.e. when $x \rightarrow -x$, if $V_1^{(1)}$ and $V_1^{(2)}$ change sign, whereas $S^{(1)}$, $S^{(2)}$, $V_0^{(1)}$ and $V_0^{(2)}$ remain the same. This is because the parity operator is $\mathcal{P} = \exp(i\delta_P) P_0 \eta^0$, where δ_P is a constant phase and P_0 changes x into $-x$. Because this unitary operator anticommutes with β^1 and $[P, \beta^1]$, they change sign under a parity transformation, whereas β^0 , P and $[P, \beta^0]$, which commute with η^0 , remain the same. Since $\delta_P = 0$ or $\delta_P = \pi$, the spinor components have definite parities and the parity of Ψ_3 is opposite to that one of Ψ_1 and Ψ_2 . The charge-conjugation operation changes the sign of the minimal interaction potential, i.e. changes the sign of $V_\mu^{(1)}$. This can be accomplished by the transformation $\Psi \rightarrow \Psi_c = \mathcal{C}\Psi = CK\Psi$, where K denotes the complex conjugation and C is a unitary matrix such that $C\beta^\mu = -\beta^\mu C$. The matrix

that satisfies this relation is $C = \exp(i\delta_C)\eta^0\eta^1$, where $\eta^1 = 2\beta^1\beta^1 + 1$. The phase factor $\exp(i\delta_C)$ is equal to ± 1 , thus $(\Psi_1)_c = \mp\Psi_1^*$, $(\Psi_2)_c = \pm\Psi_2^*$, $(\Psi_3)_c = \pm\Psi_3^*$ and $E \rightarrow -E$. Note also that $J^\mu \rightarrow -J^\mu$, as should be expected for a charge current. Meanwhile C commutes with P and anticommutes with $[P, \beta^\mu]$, then the charge-conjugation operation entails no change on $S^{(1)}$, $S^{(2)}$ and $V_\mu^{(2)}$. By the same token it can be shown that $S^{(1)}$ and $S^{(2)}$ are invariant under the time-reversal transformation and that $V_\mu^{(1)}$ and $V_\mu^{(2)}$ have opposite behavior in such a way that both sorts of vector potentials change sign under $\mathcal{PCT}$ whereas the scalar potentials do not. The invariance of $S^{(1)}$, $S^{(2)}$ and $V_\mu^{(2)}$ potentials under charge conjugation means that they do not couple to the charge of the boson. In other words, $S^{(1)}$, $S^{(2)}$ and $V_\mu^{(2)}$ do not distinguish particles from antiparticles and the spectra for those sorts of interactions are symmetrical about $E = 0$. Hence, those sorts of interactions can not exhibit Klein's paradox.

In summary, it is not correct to consider the most general form for the potential matrix as being constituted by just four Lorentz structures. There is no pseudoscalar potential in the spin-0 sector of the DKP theory. In fact, the time component of a nonminimal vector potential is used in Ref. [1]. It is true that $\beta^\mu P$ behaves like a Lorentz vector, nevertheless that term does not lead to a conserved current. The operator P considered in Ref. [1] is not the proper projection operator neither is correct the elimination $V_1^{(2)}$ by a phase transformation. Defining the transmission coefficient as the absolute value of the ratio of the transmitted flux to the incident flux one would never find out a negative transmission coefficient. It was a negative transmission coefficient obtained in Ref. [1] that allowed to conclude about the existence of Klein's paradox for a potential that does not couple to the charge of the boson. It is also worthwhile to remark that the current expressed by (3) is not a probability current but a charge current, always with $R + T = 1$ because of the conservation law. Furthermore, there is no reason to require that the spinor and its derivative are continuous across finite discontinuities of the square step potential. The proper matching conditions follow from the differential equations obeyed by the spinor components, as they should be, avoiding in this manner the hard tasking of recurring to the limit process of smooth potentials [8]. Finally, despite the restriction to the one-dimensional movement, Ref. [1] treats the problem of a particle in a (3+1)-dimensional world. In 1+1 dimensions the matrices of the DKP algebra for spin-0 particles are reduced to 3×3 matrices with three-component spinors.

Acknowledgments

This work was supported in part by means of funds provided by CAPES and CNPq.

References

- [1] A. Boumali, Can. J. Phys. **86**, 1233 (2008).
- [2] G. Petiau, Acad. R. Belg., A. Sci. Mém. Collect. **16**, No. 2 (1936).
- [3] N. Kemmer, Proc. R. Soc. A **166**, 127 (1938).
- [4] R.J. Duffin, Phys. Rev. **54**, 1114 (1938).
- [5] See, e.g., R.F. Guertin and T.L. Wilson, Phys. Rev. D **15**, 1518 (1977).
- [6] See, e.g., B. Vijayalakshmi, M. Seetharaman, and P.M. Mathews, J. Phys. A **12**, 665 (1979).
- [7] Y. Nedjadi and R.C. Barret, J. Phys. G **19**, 87 (1993).
- [8] T.R. Cardoso, L.B. Castro, and A.S. de Castro, Phys. Lett. A **372**, 5964 (2008).